

RESIDUE CURRENTS OF HOLOMORPHIC MORPHISMS

MATS ANDERSSON

ABSTRACT. Given a generically surjective holomorphic vector bundle morphism $f: E \rightarrow Q$, E and Q Hermitian bundles, we construct a current R^f with values in $\text{Hom}(Q, H)$, where H is a certain derived bundle, and with support on the set Z where f is not surjective. The main property is that if ϕ is a holomorphic section of Q , and $R^f \phi = 0$, then locally $f\psi = \phi$ has a holomorphic solution ψ . In the generic case also the converse holds. This gives a generalization of the corresponding theorem for a complete intersection, due to Dickenstein-Sessa and Passare. We also present results for polynomial mappings, related to M Noether's theorem and the effective Nullstellensatz. The construction of the current is based on a generalization of the Koszul complex. By means of this complex one can also obtain new global estimates of solutions to $f\psi = \phi$, and as an example we give new results related to the H^p -corona problem.

1. INTRODUCTION

Let E and Q be holomorphic Hermitian vector bundles of ranks m and r , respectively, over the n -dimensional complex manifold X , and let $f: E \rightarrow Q$ be a generically surjective holomorphic morphism. Given a holomorphic section ϕ of Q we are interested in holomorphic solutions ψ to $f\psi = \phi$. The basic results in this area are the existence theorems due to Skoda in [21] and [22], which are based on L^2 -methods and complex geometry. They provide existence of global holomorphic solutions to the equation $f\psi = \phi$ with L^2 -estimates under appropriate geometric conditions provided f is pointwise surjective. However, applying these results to E restricted to $X \setminus Z$, where

$$Z = \{z; f(z) \text{ is not surjective}\},$$

also highly non-trivial local results at Z are obtained by these methods.

In this paper we introduce a complex of bundles

$$\cdots \rightarrow E_3 \rightarrow E_2 \rightarrow E \rightarrow Q \rightarrow 0,$$

Date: February 2, 2008.

1991 *Mathematics Subject Classification.* 32 A 27, 32 H 02, 32 B 99.

The author was partially supported by the Swedish Natural Science Research Council.

and define a global residue current

$$R^f = R_p^f + \cdots + R_\mu^f$$

with support on Z , $p = \text{codim } Z$ and $\mu = \min(n, m - r + 1)$, where R_k^f is a $(0, k)$ -current with values in $\text{Hom}(Q, E_k)$. It is not hard to see (e.g., by using Gauss elimination) that $p \leq m - r + 1$ with equality in the generic case; in this case, thus $R = R_{m-r+1}$. Our first result concerns existence of local holomorphic solutions of $f\psi = \phi$.

Theorem 1.1. *Let E and Q be holomorphic Hermitian vector bundles over a complex manifold X , let $f: E \rightarrow Q$ be a holomorphic generically surjective morphism, and let R^f be the corresponding residue current. If ϕ is a holomorphic section of Q such that $R^f \phi = 0$, then locally $f\psi = \phi$ has a holomorphic solution ψ .*

We have the following partial converse.

Theorem 1.2. *If $p = m - r + 1$ and $f\psi = \phi$ has a holomorphic solution, then $R^f \phi = 0$.*

If $p = m - r + 1$ thus $f\psi = \phi$ has holomorphic solutions if and only if $R^f \phi = 0$. If $r = 1$ and $p = m$, it turns out that if $f = \sum f_j e_j$ in a local holomorphic frame e_j , then $R = R_m$ is equal to the classical Coleff-Herrera current

$$T = \left[\bar{\partial} \frac{1}{f_1} \wedge \cdots \wedge \bar{\partial} \frac{1}{f_m} \right]$$

times a non-vanishing section of $(\det E) \otimes Q^*$, see [4]. In this case therefore we get back the Dickenstein-Sessa-Passare theorem, [12] and [19], stating that ϕ belongs to the ideal $(f_1, \dots, f_m)$ if and only if $\phi T = 0$.

Instead of the usual norm $|\phi|$ of a section ϕ of Q it is natural, e.g., in view of the results in [22], to introduce the stronger pointwise norm

$$\|\phi\|^2 = \det(ff^*) |f^*(ff^*)^{-1} \phi|^2 = \langle \widetilde{ff^*} \phi, \phi \rangle,$$

where $\widetilde{ff^*} = \det(ff^*)(ff^*)^{-1}$ is the smooth endomorphism on Q whose matrix is the transpose of the comatrix of ff^* . By analyzing the singularity of R^f we obtain the following sufficient size condition on ϕ for annihilating the residue.

Proposition 1.3. *Let $f: E \rightarrow Q$ be a holomorphic generically surjective morphism. If ϕ is a holomorphic section of Q such that*

$$(1.1) \quad \|\phi\|^2 \leq C \det(ff^*)^{\min(n, m-r+1)},$$

then $R^f \phi = 0$.

As an immediate consequence we get the following generalization of the Briançon-Skoda theorem.

Theorem 1.4. *If ϕ is a holomorphic section of Q such that (1.1) holds, then locally $f\psi = \phi$ has a holomorphic solution ψ .*

In the case when $r = 1$, (1.1) means precisely that $|\phi| \leq C|f|^{\min(n,m)}$, and the conclusion is then that ϕ is locally in the ideal (f) generated by f (i.e., the ideal generated by f_j if $f = \sum f_j e_j$ in some local holomorphic frame e_j). This immediately implies the classical Briançon-Skoda theorem, [9], which states that $\phi^{\min(m,n)}$ belongs to (f) if $|\phi| \leq C|f|$. For $m - r + 1 \leq n$, Theorem 1.3 also follows directly from Skoda's L^2 -estimate, [22], but when $m - r + 1 > n$, the L^2 -estimate only gives the conclusion if the power of the right hand side of (1.1) is $n + 1$. By an additional argument in the case when $m > n$, the classical Briançon-Skoda theorem follows from the L^2 -estimate; the case when $m > n$ and $r > 1$ can be reduced to the classical result by means of the Fuhrmann trick, see Section 7.

Demailly has extended Skoda's L^2 -theorems to $\bar{\partial}$ -closed sections, see [10] and [11]. Our method also admits such an extension of the local result.

Theorem 1.5. *Assume that ϕ is a smooth $\bar{\partial}$ -closed $(0, q)$ -section of Q . If $R^f \phi = 0$, then locally $f\psi = \phi$ has $\bar{\partial}$ -closed (current) solutions ψ .*

For degree reasons we see that $R^f \phi = 0$ if $q > n - p$. Thus we get

Corollary 1.6. *If ϕ is any smooth $\bar{\partial}$ -closed $(0, q)$ -form with values in Q , and $q > n - p = \dim Z$, then locally $f\psi = \phi$ has $\bar{\partial}$ -closed current solutions.*

In analogy with Theorems 1.3 and (1.4) we also have

Theorem 1.7. *If ϕ is a smooth $\bar{\partial}$ -closed $(0, q)$ -form with values in Q such that*

$$(1.2) \quad \|\phi\|^2 \leq C \det(ff^*)^{\min(n-q, m-1+r)},$$

then $R^f \phi = 0$, and locally there are integrable $\bar{\partial}$ -closed solutions ψ to $f\psi = \phi$.

We can also obtain global results and first we turn our attention to polynomial ideals and generalize the approach in [5]. Let $[z] = [z_0, \dots, z_n]$ be homogeneous coordinates on $\mathbb{P}^n$, and let $z' = (z_1, z_2, \dots, z_n)$ be the standard coordinates in the standard affinization $\mathbb{C}^n \simeq \{[z], z_0 \neq 0\}$. Let P be a polynomial mapping $\mathbb{C}^n \rightarrow \text{Hom}(\mathbb{C}^m, \mathbb{C}^r)$ with columns P^j such that $\deg P^j \leq d_j$, $j = 1, \dots, m$. If f is the matrix whose columns are the d_j -homogenized forms $f^j(z) = z_0^{d_j} P^j(z'/z_0)$ in $\mathbb{C}^{n+1}$, then f defines a morphism

$$f: \bigoplus_{j=1}^m \mathcal{O}(-d_j) \rightarrow \mathbb{C}^r.$$

Let Z be the algebraic variety in $\mathbb{P}^n$ where f is not surjective, and let R^f be the associated residue current with respect to the natural metric.

Theorem 1.8. *Assume that P is a polynomial mapping as above, and let Φ be a r -column of polynomials of degrees $\leq \rho$. Moreover, assume that*

$$(1.3) \quad m \leq n + r - 1 \quad \text{or} \quad \rho \geq \sum_{j=1}^{n+r} d_j - n,$$

where $d_1 \geq d_2 \geq \dots \geq d_m$. If $R^f \phi = 0$, then there are polynomials Q_j such that $\sum_1^m P^j Q_j = \Phi$, and $\deg P^j Q_j \leq \rho$.

Corollary 1.9. *Assume that Z is empty. Then we can find a matrix Q of polynomials with rows Q_k such that $PQ = \sum P^k Q_k = I_r$, and $\deg P^k Q_k \leq \sum_{j=1}^{n+r} d_j - n$.*

This is a generalization of a classical theorem of Macaulay, [17].

Corollary 1.10. *Let Φ be a column of polynomials, $\deg \Phi \leq \rho$, and let ϕ be its ρ -homogenization. If*

$$\|\phi\|^2 \leq C \det(f f^*)^{\min(n, m-r+1)}$$

in $\mathbb{P}^n$, and (1.3) is fulfilled, then $PQ = \Phi$ has a solution with $\deg P^k Q_k \leq \rho$.

Assume that P is pointwise surjective in $\mathbb{C}^n$, and let $P^j = (P_1^j, \dots, P_r^j)^t$. By the local Lojasiewicz inequality there is a constant M such that

$$(1.4) \quad \sum_{|I|=r} \frac{|\det(P_k^{I_j}(z'))|^2}{(1 + |z'|^2)^{\sum_1^r d_{I_j}}} \geq C \frac{1}{(1 + |z'|^2)^M},$$

where the sum is over increasing multiindices.

Corollary 1.11. *Assume that $P: \mathbb{C}^n \rightarrow \text{Hom}(\mathbb{C}^m, \mathbb{C}^r)$ is surjective in $\mathbb{C}^n$, $\deg P \leq d$ and that (1.4) holds. Then there is a matrix Q of polynomials such that $PQ = I_r$ and $\deg P^j Q_j \leq M \min(n, m - r + 1)$.*

From Kollar's famous theorem, [16], we can get an estimate of M . For simplicity we assume $d_j = d$ for all j .

Proposition 1.12. *If $d_j = d$ for all j , then the inequality (1.4) holds with*

$$M = (rd)^{\min(n, m!/(m-r)!r!)},$$

provided that $rd \geq 3$.

It should be pointed out that the bound

$$\deg Q + d \leq \min(n, m - r + 1)M$$

we obtain in this way for a solution to $PQ = I_r$ (even) is not optimal when $r = 1$. It is proved in [16] that one actually have $\deg Q + d \leq M$

when $r = 1$. We do not know if it is possible to modify Kollar's proof as to include the case $r > 1$ directly and get a sharper bound.

Now assume that $p = m - r + 1 = n$ and that Z is contained in $\mathbb{C}^n$; thus a discrete set. Moreover, assume that $\Phi = PQ$ is solvable in $\mathbb{C}^n$. Then it follows from Theorem 1.2 that $R^f\phi = 0$ in $\mathbb{C}^n$, and hence $R^f\phi = 0$ in $\mathbb{P}^n$, and since (1.3) is fulfilled we can take $\rho = \deg \Phi$. Therefore there is a solution to $PQ = \Phi$ such that $\deg P^j Q_j \leq \deg \Phi$. When $r = 1$ this is a classical theorem due to Max Noether, [18]. We have the following generalization that appeared in [5] in the case $r = 1$; however we suspect that this case could be proved algebraically, e.g., by the methods in [15].

Theorem 1.13. *Assume that $P: \mathbb{C}^n \rightarrow \text{Hom}(\mathbb{C}^m, \mathbb{C}^r)$ and that $p = m - r + 1$ and that Z has no irreducible component contained in the hyperplane at infinity. Moreover, assume that $\Phi = PQ$ is solvable in $\mathbb{C}^n$. Then there is a solution Q such that $\deg P^j Q_j \leq \deg \Phi$.*

We can also obtain new global results in open bounded domains even when f is pointwise surjective, and as an example we present in Section 7 new sharpened estimates of solutions to the H^p -corona problem in a strictly pseudoconvex domain.

CONTENTS

1. Introduction	1
2. A generalized Koszul complex	5
3. Surjective morphisms	8
4. Analysis of singularities	10
5. The residue current of a generically surjective holomorphic morphism	13
6. Polynomial mappings	16
7. Estimates for a pointwise surjective morphism	17
References	20

2. A GENERALIZED KOSZUL COMPLEX

Let $f: E \rightarrow Q$ be a holomorphic morphism as above. Assume that we have a complex

$$(2.1) \quad \cdots \rightarrow E_3 \rightarrow E_2 \rightarrow E \rightarrow Q \rightarrow 0.$$

of vector bundles where all the morphisms, which we denote by δ , are holomorphic. Let $E_0 = Q$, $E_1 = E$, and let

$$H = \bigoplus_{k=0}^{\infty} E_k.$$

We will also consider $(0, *)$ -form-valued sections of H , i.e., sections of $T_{0,1}^*(X) \otimes H$. We denote this space of sections by $\mathcal{E}_{0,*}(X, H)$. Notice

that it is a module over the ring (algebra) $\mathcal{E}_{0,*}(X)$. We extend the action of δ to sections of $T_{0,1}^*(X) \otimes H$ by requiring that

$$(2.2) \quad \delta\xi \otimes w = (-1)^{\deg \xi} \xi \otimes \delta w$$

if ξ is a differential form. Then $\delta\bar{\partial} = -\bar{\partial}\delta$.

Now suppose that we have $(0, k-1)$ -forms, or currents, v_k with values in E_k , $k \geq 1$, such that

$$(2.3) \quad (\delta - \bar{\partial})(v_1 + v_2 + \cdots) = \phi,$$

i.e.,

$$(2.4) \quad \delta v_{k+1} = \bar{\partial} v_k, \quad k \geq 1, \quad \delta v_1 (= f v_1) = \phi.$$

For degree reasons, $\bar{\partial} v_k = 0$ if k is large enough, and if there are no obstructions for solving $\bar{\partial}$, we can successively find $(0, k-2)$ -forms (currents) w_k with values in E_k such that

$$(2.5) \quad \bar{\partial} w_k = v_k + \delta w_{k+1}, \quad k \geq 2.$$

Then finally

$$(2.6) \quad \psi = v_1 + \delta w_2$$

is a holomorphic solution to $f\psi = \phi$. Since the $\bar{\partial}$ -equations always are solvable locally we have

Lemma 2.1. *Suppose that we have a current solution $v = v_1 + v_2 + \dots$ to (2.3). Then locally there are holomorphic solutions to $f\psi = \phi$.*

If f is surjective, then obviously there are local holomorphic solutions to $f\psi = \phi$ so the interesting case is when f is just generically surjective. In view of the argument in the proof, one also gets a global holomorphic solution provided all the $\bar{\partial}$ -equations have global solutions. Before we proceed with our construction let us consider some examples.

Example 1. If the complex (2.1) is exact (in particular f is surjective), then we can always find such a solution to (2.3). In fact, given a holomorphic section ϕ of Q , let v_1 be any pointwise solution to $\delta v = \phi$. Then $\bar{\partial} v_1$ is δ -exact and hence there is a v_2 such that $\delta v_2 = \bar{\partial} v_1$ etc. $\square$

Example 2. If $r = 1$, i.e., Q is a line bundle, then one can take $E_k = \Lambda^k E \otimes Q^*$ and δ as interior multiplication with f . One then gets the usual Koszul complex

$$(2.7) \quad \cdots \rightarrow \Lambda^3 E \otimes (Q^*)^2 \xrightarrow{\delta} \Lambda^2 E \otimes Q^* \xrightarrow{\delta} E \xrightarrow{f} Q \rightarrow 0,$$

which is exact if (and only if) f is non-vanishing. In fact, if we choose any section of $\text{Hom}(Q, E) \simeq E \otimes Q^*$ $f\sigma = I_Q$ (if E has a Hermitian metric we can, e.g., choose the section with pointwise minimal norm), then there is an induced mapping $\sigma: \Lambda^k E \otimes (Q^*)^{k-1} \rightarrow \Lambda^{k+1} E \otimes (Q^*)^k$ such that $\delta \circ \sigma + \sigma \circ \delta = I$, and thus (2.7) is exact. $\square$

Example 3. Provided that f is surjective, a simple way to find an exact complex (2.1) is by taking $E_2 = \text{Ker } f$ and $E_k = 0$ for $k > 2$. However, E_2 is usually not trivial in a neighborhood of a singular point, i.e., E_2 usually cannot be extended as a vector bundle across the set Z where f is not surjective. To carry out the scheme in the proof of Lemma 2.1, one therefore has to solve a $\bar{\partial}$ -equation in the bundle $E_2 = \text{Ker } f$ over $X \setminus Z$. This is only possible under certain geometric conditions; this is precisely what is investigated and explained in [22]. $\square$

Let us now describe our generalized Koszul complex. Notice that our holomorphic morphism $f: E \rightarrow Q$ is a holomorphic section of the bundle $\text{Hom}(E, Q)$, which we identify with $E^* \otimes Q$. If ϵ_j is a local holomorphic frame for Q , then

$$f = \sum f_j \otimes \epsilon_j,$$

where f_j are sections of E^* . If η is a section of E , then $f\eta = \sum_1^r \delta_{f_j} \epsilon_j$, where δ_{f_j} denotes interior multiplication with f_j . We can associate to f the section

$$F = f_1 \wedge f_2 \dots \wedge f_r \otimes \epsilon_1 \wedge \dots \wedge \epsilon_r$$

of $\Lambda^r E^* \otimes \det Q^*$. It is independent of the particular choice of frame, and will be called the determinant section of f . Notice that f is surjective at a point if and only if F is non-vanishing at that point. There is an induced mapping

$$\delta_F: \Lambda^{r+1} E \otimes \det Q^* \rightarrow E$$

defined by

$$\delta_F(\xi \otimes \epsilon_1^* \wedge \dots \wedge \epsilon_r^*) = \delta_{f_r} \dots \delta_{f_1} \xi,$$

which is also easily seen to be independent of the particular local frame ϵ_j for Q ; here ϵ_j^* denotes the dual frame for Q^* . Moreover, it is also clear that

$$(2.8) \quad f \circ \delta_F = 0.$$

In order to proceed with the construction of our complex we have to recall some facts about symmetric tensors. Let $S^\ell Q^*$ be the subbundle of $\bigotimes Q^*$ consisting of symmetric ℓ -tensors of Q^* . If $u, v \in Q^*$ then $u \dot{\otimes} v = u \otimes v + v \otimes u$, etc. This extends to a commutative mapping $SQ^* \times SQ^* \rightarrow SQ^*$. If q is a section of Q , then it induces the usual interior multiplication on $\bigotimes Q^*$ (say from the left), and, in particular, if $u^p = u \dot{\otimes} \dots \dot{\otimes} u$, then $\delta_q u^p = p u^{p-1} (q \cdot u)$.

We now define

$$(2.9) \quad E_k = \Lambda^{r+k-1} E \otimes S^{k-2} Q^* \otimes \det Q^*, \quad k \geq 2.$$

Given the local frame ϵ_j , a section ξ of E_k can be written

$$\xi = \sum_{|\alpha|=k-2} \xi_\alpha \otimes \epsilon_\alpha^* \otimes \epsilon^*,$$

where

$$\epsilon_\alpha^* = \frac{(\epsilon_1^*)^{\alpha_1} \dot{\otimes} \dots \dot{\otimes} (\epsilon_r^*)^{\alpha_r}}{\alpha_1! \dots \alpha_r!}$$

and $\epsilon^* = \epsilon_1^* \wedge \dots \wedge \epsilon_r^*$. For $k \geq 2$ we have mappings $\delta: E_{k+1} \rightarrow E_k$ defined by

$$\xi \otimes q^* \otimes \epsilon^* \mapsto \sum_{j=1}^r \delta_{f_j} \xi \otimes \delta_{\epsilon_j} q^* \otimes \epsilon^*,$$

which are also independent of the specific choice of local frame ϵ_j . Since δ_{f_j} anti-commute and δ_{ϵ_j} commute, it follows that $\delta^2 = 0$. Moreover, if the section ξ of E_2 is in the image of δ , i.e., $\xi = \delta\eta = \sum \delta_{f_j} \eta_j \otimes \epsilon^*$, then clearly $\delta_F \xi = 0$. In view of (2.8) we thus have a complex

$$(2.10) \quad \dots \xrightarrow{\delta} E_3 \xrightarrow{\delta} E_2 \xrightarrow{\delta_F} E \xrightarrow{f} Q \rightarrow 0.$$

In the sequel, we will often denote all the mappings in (2.10) by δ . Observe that if $r = 1$, then (2.10) is just the Koszul complex (2.7).

If we let ξ above take values in $\Lambda(T^*(X)_{0,1} \oplus E)$ rather than just ΛE , then we get an extension of all the mappings δ and δ_F to forms and currents with values in E_k . The mappings $\delta: E_{k+1} \rightarrow E_k$, $k \geq 2$, will automatically satisfy (2.2) so that $\delta\bar{\partial} = -\bar{\partial}\delta$, but we should have to insert the factor $(-1)^{(r+1)q}$ in the definition of δ_F , when it acts on $\xi \otimes w$, and ξ is a $(0, q)$ -form. However, we will not do that, and therefore we have instead that $\bar{\partial}\delta_F = (-1)^r \delta_F \bar{\partial}$. This means that the final solution ψ in (2.6) is $\psi = v_1 + (-1)^{r+1} \delta w_2$.

3. SURJECTIVE MORPHISMS

Now we assume that $f: E \rightarrow Q$ is surjective and that E and Q are equipped with Hermitian metrics. Moreover, we let E_k be the derived bundles defined by (2.9). Let σ be the section of $\text{Hom}(Q, E) = E \otimes Q^*$ with pointwise minimal norm (i.e., such that $\text{Im } \sigma$ is orthogonal to $\text{Ker } f$) such that $f \circ \sigma = I_Q$. If ϵ_j^* denotes the dual frame for Q^* , then

$$\sigma = \sum \sigma_j \otimes \epsilon_j^*,$$

where σ_j are the sections of E with minimal norms such that $f_j \cdot \sigma_k = \delta_{jk}$. Moreover,

$$\sigma = \sigma_1 \wedge \dots \wedge \sigma_r \otimes \epsilon_1^* \wedge \dots \wedge \epsilon_r^*$$

is a well-defined section of $\Lambda E^r \otimes \det Q^*$, and it induces a mapping $E \rightarrow E_2 = \Lambda^{r+1} E \otimes \det Q^*$ defined by

$$\xi \mapsto \sigma \wedge \xi = \sigma_1 \wedge \dots \wedge \sigma_r \wedge \xi \otimes \epsilon_1^* \wedge \dots \wedge \epsilon_r^*.$$

Now, $\delta_F \sigma \xi = \xi$ provided that $\delta_{f_j} \xi = 0$ for all j , i.e., ξ is in $\text{Ker } f$. Thus (2.10) is exact at E if f is surjective. We also have

Lemma 3.1. *If f is surjective, then (2.10) is exact up to E_2 .*

Proof. It remains to check the exactness at E_2 . Suppose that $\xi \otimes \epsilon^*$ is a section of E_2 such that $0 = \delta_F \xi \otimes \epsilon^*$. Thus ξ is a section of $\Lambda^{r+1}E$, such that $\delta_{f_r} \cdots \delta_{f_1} \xi = 0$. By the surjectivity of f , σ_j are linearly independent and therefore they form a part of a basis $\sigma_1, \dots, \sigma_m$ for E . With respect to this basis we can write

$$\xi = \sigma_1 \wedge \dots \wedge \sigma_r \wedge \xi' + \xi'',$$

where ξ'' does not contain all the σ_j , $j = 1, \dots, r$. It follows that $\xi' = 0$ and hence $\xi = \sum_1^r \xi_j$, where ξ_j does not contain σ_j . Therefore

$$\xi \otimes \epsilon^* = \sum_1^r \delta_{f_j}(\sigma_j \wedge \xi_j) \otimes \epsilon^* = \delta\left(\sum_1^r \sigma_j \wedge \xi_j \otimes \epsilon_j^* \otimes \epsilon^*\right),$$

and thus (2.10) is exact at E_2 . $\square$

In order to find a solution to (2.3) it is natural to start with $v_1 = \sigma\phi = \sum \phi_j \sigma_j$, and $v_2 = \sigma \wedge \bar{\partial}\sigma\phi$. We can just as well suppress ϕ and define u_k with values in $\text{Hom}(Q, E_k)$ such that $u_k\phi$ satisfies (2.3). Notice that $\text{Hom}(Q, E_1) \simeq E \otimes Q^*$ and

$$\text{Hom}(Q, E_k) \simeq \Lambda^{r+k-1}E \otimes S^{k-2}Q^* \otimes \det Q^* \otimes Q^*, \quad k \geq 2.$$

Definition 1. We define the $(0, k-1)$ -forms u_k with values in $\text{Hom}(Q, E_k)$ as

$$(3.1) \quad u_1 = \sigma, \quad u_k = (\bar{\partial}\sigma)^{\otimes(k-2)} \otimes \sigma \otimes \bar{\partial}\sigma, \quad k \geq 2.$$

Notice that $(\bar{\partial}\sigma)^{\otimes(k-2)}$ is indeed a symmetric tensor, and that the definition of u_k is invariant. Moreover,

$$\left(\sum_1^r \bar{\partial}\sigma_\ell \otimes \epsilon_\ell^*\right)^{\otimes(k-2)} = \frac{(\sum_1^r \bar{\partial}\sigma_\ell \otimes \epsilon_\ell^*)^{\dot{\otimes}(k-2)}}{(k-2)!} = \sum_{|\alpha|=k-2} (\bar{\partial}\sigma)^\alpha \otimes \epsilon_\alpha^*,$$

where

$$(\bar{\partial}\sigma)^\alpha = (\bar{\partial}\sigma_1)^{\alpha_1} \wedge \dots \wedge (\bar{\partial}\sigma_r)^{\alpha_r}.$$

Since $\bar{\partial}\sigma_j$ have degree 2 in $\Lambda(T_{0,1}^*(X) \oplus E)$ and therefore commute, we thus have that

$$(3.2) \quad u_1 = \sigma = \sum_j \sigma_j \otimes \epsilon_j^*,$$

$$u_k = \sigma_1 \wedge \dots \wedge \sigma_r \wedge \sum_{|\alpha|=k-2} \sum_j (\bar{\partial}\sigma)^\alpha \wedge \bar{\partial}\sigma_j \otimes \epsilon_\alpha^* \otimes \epsilon^* \otimes \epsilon_j^*, \quad k \geq 2.$$

Proposition 3.2. Assume that f is surjective. If u is defined by (3.1), then

$$(\delta - \bar{\partial})(u_1 + u_2 + \dots) = I_Q.$$

Here $I_Q: Q \rightarrow Q$ is the identity morphism.

Proof. We have already seen that $\delta u_1 = f\sigma = I_Q$ and $\delta_F \sigma \otimes \bar{\partial}\sigma = \bar{\partial}\sigma = \bar{\partial}u_1$, so we have to verify that $\delta u_{k+1} = \bar{\partial}u_k$ for $k \geq 2$. Now,

$$u_{k+1} = \sigma_1 \wedge \dots \wedge \sigma_r \wedge \left(\sum_1^r \bar{\partial}\sigma_\ell \otimes \epsilon_\ell^* \right)^{\otimes(k-1)} \otimes \epsilon^* \otimes \bar{\partial}\sigma.$$

Recalling that $\delta = \sum \delta_{f_j} \otimes \delta_{\epsilon_j}$, and that δ_{ϵ_j} acts from the left, we get

$$\delta u_{k+1} = \sum_{j=1}^r \delta_{f_j} (\sigma_1 \wedge \dots \wedge \sigma_r) \wedge \bar{\partial}\sigma_j \wedge \left(\sum_1^r \bar{\partial}\sigma_\ell \otimes \epsilon_\ell^* \right)^{\otimes(k-2)} \otimes \epsilon^* \otimes \bar{\partial}\sigma.$$

On the other hand,

$$\bar{\partial}u_k = \bar{\partial}(\sigma_1 \wedge \dots \wedge \sigma_r) \wedge \left(\sum_1^r \bar{\partial}\sigma_\ell \otimes \epsilon_\ell^* \right)^{\otimes(k-2)} \otimes \epsilon^* \otimes \bar{\partial}\sigma.$$

It is now clear that $\delta u_{k+1} = \bar{\partial}u_k$. $\square$

It follows that if $\phi = \sum \phi_j \epsilon_j$ is a holomorphic section of Q and $v_k = u_k \phi$, then $v = u\phi$ satisfies (2.3). For later purpose we rewrite the expression for $u\phi$ so that it only “depends” on $\sigma\phi = \sum_j \phi_j \sigma_j$. In fact, applying $\bar{\partial}$ to the equality $0 = \sigma \otimes \sigma$ we get $0 = \sigma \otimes \bar{\partial}\sigma + (-1)^r \bar{\partial}\sigma \otimes \sigma$. Therefore,

$$(3.3) \quad u_k \phi = (-1)^{r+1} (\bar{\partial}\sigma)^{\otimes(k-2)} \otimes \bar{\partial}\sigma \otimes \sigma\phi, \quad k \geq 2,$$

or more explicitly (the possible minus sign cancels out)

$$\begin{aligned} u_k \phi &= \\ & \left(\sum_j \phi_j \sigma_j \right) \wedge \bar{\partial}(\sigma_1 \wedge \dots \wedge \sigma_r) \wedge \sum_{|\alpha|=k-2} (\bar{\partial}\sigma)^\alpha \otimes \epsilon_\alpha^* \otimes \epsilon^* = \\ & \left(\sum_j \phi_j \sigma_j \right) \wedge \bar{\partial}(\sigma_1 \wedge \dots \wedge \sigma_r) \wedge \left(\sum_1^r \bar{\partial}\sigma_\ell \otimes \epsilon_\ell^* \right)^{\otimes(k-2)} \otimes \epsilon^*, \quad k \geq 2. \end{aligned}$$

4. ANALYSIS OF SINGULARITIES

We now consider an f that is not necessarily surjective everywhere. Then we can define u as above in $X \setminus Z$, where Z is the set where f is not surjective, which is equal to the zero set of the holomorphic section F and hence an analytic subvariety of X . To analyze the singularities of u at Z we will use the following lemma.

Lemma 4.1. (i) Let s be the section of $\text{Hom}(Q, E) \simeq E \otimes Q^*$ with pointwise minimal norm such that

$$fs = |F|^2 I_Q,$$

and let S be the section of $\Lambda^r E \otimes \det Q^*$ with pointwise minimal norm such that

$$FS = 1.$$

Then s and S are smooth across Z , and

$$s = |F|^2 \sigma, \quad S = |F|^2 \sigma \quad \text{in } X \setminus Z.$$

(ii) If in addition $F = F_0 F'$ for some holomorphic function F_0 and non-vanishing holomorphic section F' , then

$$s' = F_0 \sigma, \quad S' = F_0 \sigma$$

are smooth across Z .

Given a section η of E^* , let η^* denote the dual section with respect to the Hermitian metric, i.e., $\langle \xi, \eta^* \rangle = \eta \cdot \xi$ for sections ξ of E . The mapping $\eta \mapsto \eta^*$ is conjugate-linear and extends to a conjugate-linear mapping $\Lambda E^* \rightarrow \Lambda E$ by

$$\eta_1 \wedge \dots \wedge \eta_r \mapsto \eta_1^* \wedge \dots \wedge \eta_r^*.$$

Proof. Assume that ϵ_j is a local frame for Q as before, let ϵ_j^* be its dual frame, and assume $f = \sum f_j \otimes \epsilon_j$. Now,

$$S = |\epsilon_1 \wedge \dots \wedge \epsilon_r|^2 f_1^* \wedge \dots \wedge f_r^* \otimes \epsilon_1^* \wedge \dots \wedge \epsilon_r^*$$

is a section of $\Lambda^r E \otimes \det Q^*$ that is independent of the particular choice of frame. This is checked by considering a change of frame $\epsilon' = \epsilon g$, where g is an invertible $r \times r$ -matrix. Notice that

$$(4.1) \quad FS = |\epsilon_1 \wedge \dots \wedge \epsilon_r|^2 |f_1 \wedge \dots \wedge f_r|^2 = |F|^2.$$

We can choose the frame ϵ_j such that f_j are orthogonal at any given point outside Z . Thus $f_j = \alpha_j e_j^*$, $j = 1, \dots, r$ for some ON -frame e_j^* of E^* and $f_j^* = \bar{\alpha}_j e_j$, and it is then easy to see that S is in fact the section with minimal norm such that $FS = 1$. In particular this means that S is the dual section of F . Moreover, at this point, $\sigma_j = (1/\alpha_j) e_j$, $j = 1, \dots, r$, and thus

$$\sigma = \frac{e_1 \wedge \dots \wedge e_r}{\alpha_1 \dots \alpha_r} \otimes \epsilon_1^* \wedge \dots \wedge \epsilon_r^*.$$

Therefore,

$$|F|^2 \sigma = |\epsilon_1 \wedge \dots \wedge \epsilon_r|^2 \bar{\alpha}_1 \dots \bar{\alpha}_r e_1 \wedge \dots \wedge e_r \otimes \epsilon_1^* \wedge \dots \wedge \epsilon_r^* = S.$$

Now assume that $F = F_0 F'$. Since S is the dual section of F it follows that $S = \bar{F}_0 S''$, where S'' is the dual section of F' , and thus S'' is smooth even across Z . Therefore $F_0 \sigma = F_0 S / |F|^2 = S'' / |F'|^2$ is smooth across Z as well.

Now, define

$$s = \left(\sum_1^r \delta_{f_j} \otimes \delta_{\epsilon_j} \right)^{r-1} S / r! =$$

$$|\epsilon_1 \wedge \dots \wedge \epsilon_r|^2 \sum_1^r (-1)^{\ell+1} \delta_{f_r} \dots \delta_{f_{\ell+1}} \delta_{f_{\ell-1}} \dots \delta_{f_1} (f_1^* \wedge \dots \wedge f_r^*) \otimes \epsilon_j^*.$$

Clearly,

$$fs = \sum_1^r \delta_{f_j} \otimes \epsilon_j s = |\epsilon_1 \wedge \dots \wedge \epsilon_r|^2 |f_1 \wedge \dots \wedge f_r|^2 \sum_j \epsilon_j \otimes \epsilon_j^* = |F|^2 I_Q.$$

Moreover, it is readily checked that s is orthogonal to $\text{Ker } f$ so that s is the minimal section such that $fs = |F|^2 I_Q$. One can also check this by choosing a frame as above such that that f_j are orthogonal. Thus, $s/|f|^2 = \sigma$. Finally, if $F_0 \sigma$ is smooth across Z it follows that

$$F_0 \sigma = \left(\sum_1^r \delta_{f_j} \otimes \delta_{\epsilon_j} \right)^{r-1} F_0 \sigma$$

is smooth as well. $\square$

Remark 1. Assume that Q as well as E are trivial and equipped with the trivial metrics, and e_j and ϵ_j are ON-frames. If

$$F = \sum_{|I|=r}^I F_I \otimes e_I^*,$$

(suppressing the factor $\epsilon_1 \wedge \dots \wedge \epsilon_r$ and its dual), we have that

$$S = \sum_{|I|=r}^I \overline{F_I} \otimes e_I.$$

$\square$

Since $\sigma \otimes \sigma = 0$ we have, for any scalar function ξ , that

$$(4.2) \quad (\bar{\partial}(\xi\sigma))^{\otimes(k-2)} \otimes \xi\sigma \otimes \bar{\partial}(\xi\sigma) = (-1)^{r+1} (\bar{\partial}(\xi\sigma))^{\otimes(k-2)} \otimes \bar{\partial}(\xi\sigma) \otimes \xi\sigma = \xi^k u_k.$$

In view of Lemma 4.1 we have in particular that

$$(4.3) \quad u_1 = \frac{s}{|F|^2},$$

$$u_k = \frac{(\bar{\partial}s)^{\otimes(k-2)} \otimes S \otimes \bar{\partial}s}{|F|^{2k}} = (-1)^{r+1} \frac{(\bar{\partial}s)^{\otimes(k-2)} \otimes \bar{\partial}S \otimes s}{|F|^{2k}}, \quad k \geq 2$$

Lemma 4.2. *If $f^*: Q \rightarrow E$ is the adjoint morphism with respect to the Hermitian structures on Q and E , then*

$$(4.4) \quad |F|^2 = \det(ff^*).$$

Proof. Since both sides of (4.4) are invariant pointwise statements, we can assume that ϵ_j is a ON-frame. Let ξ be any section of E . Then

$$f_\ell \cdot \xi = \left\langle \sum_j (f_j \cdot \xi) \epsilon_j, \epsilon_\ell \right\rangle = \langle f\xi, \epsilon_\ell \rangle = \langle \xi, f^* \epsilon_\ell \rangle$$

which means that $f^* = \sum_j f_j^* \otimes \epsilon_j^*$. It follows that

$$ff^* = \sum (f_j \cdot f_k^*) \epsilon_j \otimes \epsilon_k^* = \langle f_j, f_k \rangle \epsilon_j \otimes \epsilon_k^*.$$

Thus

$$\det(ff^*) = \det \langle f_j, f_k \rangle = |f_1 \wedge \dots \wedge f_r|^2,$$

which implies the statement since $|\epsilon_1 \wedge \dots \wedge \epsilon_r| = 1$, cf., (4.1). $\square$

Since $(ff^*)^{-1} = \widetilde{ff^*} / \det(ff^*)$ and $s/|F|^2 = \sigma = f^*(ff^*)^{-1}$ we can conclude that

$$s = \widetilde{f^* f f^*}.$$

Moreover, if $F = F_0 F'$ as in Lemma 4.1 above, then by (4.4),

$$(4.5) \quad |s' \phi| = |\sigma \phi| |F_0| = |f^*(ff^*)^{-1} \phi| |F|/|F'| = \|\phi\|/|F'|.$$

5. THE RESIDUE CURRENT OF A GENERICALLY SURJECTIVE HOLOMORPHIC MORPHISM

We say that $f: E \rightarrow Q$ is generically surjective if Z has positive codimension. Again let E_k be the derived bundles defined by (2.9) and let $H = Q \oplus E \oplus E_2 \oplus E_3 \oplus \dots$. From Section 3 we have the section u of $\text{Hom}(Q, H)$ over $X \setminus Z$ defined by (3.2). Following [4] we shall now extend it to a current U with values in $\text{Hom}(Q, H)$ across Z and define the corresponding residue current.

Theorem 5.1. *Assume that $f: E \rightarrow Q$ is a generically surjective holomorphic morphism and let u be the associated section of $\text{Hom}(Q, H)$ defined in $X \setminus Z$. The function $\lambda \mapsto |F|^{2\lambda} u$ is holomorphic for $\text{Re } \lambda > -\epsilon$ and*

$$U = |F|^{2\lambda} u|_{\lambda=0}$$

is a current extension of u across Z . Moreover,

$$(\delta - \bar{\partial})U = I_Q - R^f,$$

where

$$R^f = \bar{\partial}|F|^{2\lambda} \wedge u|_{\lambda=0}.$$

The current R^f (taking values in $\text{Hom}(Q, H)$) has support on Z and

$$R^f = R_p^f + \dots + R_\mu^f,$$

where $p = \text{codim } Z$ and $\mu = \min(n, m - r + 1)$, and R_k^f is a current of bidegree $(0, k)$ with values in E_k .

Proof. The proof is more or less identical to the proof of Theorem 1.1 in [4], so we only sketch it. After an appropriate resolution of singularities we may assume that $F = F_0 F'$, where F_0 is a holomorphic function and F' is a non-vanishing section of (the pullback of) $\Lambda^r E^* \otimes \det Q$.

According to Lemma 4.1 (ii), then $s' = F_0\sigma$ and $S' = F_0\sigma$ are smooth across the singularity, and hence by (4.2),

$$u_1 = \frac{s'}{F_0}, \quad u_k = (-1)^{r+1} \frac{(\bar{\partial}s')^{\otimes(k-2)} \otimes \bar{\partial}S' \otimes s'}{F_0^k}, \quad k \geq 2.$$

It is now easy to see that the analytic continuations of $|F|^{2\lambda}u_k$ exist, and in this resolution the values at $\lambda = 0$ are just the principal value currents

$$u_1 = \left[\frac{1}{F_0} \right] s', \quad \left[\frac{1}{F_0^k} \right] (\bar{\partial}s')^{\otimes(k-2)} \otimes \bar{\partial}S' \otimes s'.$$

Precisely as in [4], by the way following [7] and [20], one can show that

$$R_k^f = \bar{\partial}|F|^{2\lambda} \wedge u_k|_{\lambda=0} = 0$$

if $k < p = \text{codim } Z$. Thus Theorem 5.1 is proved. $\square$

Proof of Theorem 1.2. If

$$\phi = f\psi = \sum_1^r (\delta_{f_j}\psi)\epsilon_j,$$

then

$$\begin{aligned} u_{m-r+1}\phi &= \left(\sum_1^r \phi_j \bar{\partial}\sigma_j \right) \wedge \sigma_1 \wedge \dots \wedge \sigma_r \wedge (\bar{\partial}\sigma)^{\otimes(m-r-1)} \otimes \epsilon^* = \\ &= \left(\sum_{j=1}^r \delta_{f_j}\psi \wedge \bar{\partial}\sigma_j \right) \wedge \sigma_1 \wedge \dots \wedge \sigma_r \wedge (\bar{\partial}\sigma)^{\otimes(m-r-1)} \otimes \epsilon^* = \\ &= \psi \wedge \sum_{j=1}^r \delta_{f_j}(\sigma_1 \wedge \dots \wedge \sigma_r) \wedge \bar{\partial}\sigma_j \wedge (\bar{\partial}\sigma)^{\otimes(m-r-1)} \otimes \epsilon^* = \\ &= \bar{\partial}(\psi \wedge \sigma_1 \wedge \dots \wedge \sigma_r \wedge (\bar{\partial}\sigma)^{\otimes(m-r-1)} \otimes \epsilon^*) = \bar{\partial}(u'\psi), \end{aligned}$$

where we have used that the form has maximal degree m in ΛE . If we define

$$R'\psi = \bar{\partial}|f|^{2\lambda} \wedge u' \wedge \psi|_{\lambda=0},$$

it follows that

$$R_{m-r+1}\phi = \bar{\partial}(R'\psi).$$

However, since $\text{codim } Z = m - r + 1$ it follows as in the proof of Theorem 5.1 above that $R'\psi$, being a $(0, m - r)$ -current, vanishes for degree reasons. Thus the theorem is proved. $\square$

Proof of Proposition 1.3. Let ξ be a test form with support contained in neighborhood where we have the resolution of singularities. In view of (4.2) we see that $R_k^f \cdot \xi$ is a sum of terms like

$$\bar{\partial}|F_0|^{2\lambda} v^\lambda \wedge \frac{(\bar{\partial}s')^{\otimes(k-2)} \otimes \bar{\partial}S' \otimes s' \phi}{F_0^k} \rho \Big|_{\lambda=0},$$

where ρ is a cut-off function, and v is a smooth strictly positive function. By the hypothesis, (4.5), and (4.2),

$$|s'\phi| \sim \|\phi\| \lesssim |F|^{\min(n, m-r+1)} \lesssim |F_0|^k,$$

since $k \leq \min(n, m-r+1)$. Thus we must check that

$$\int \bar{\partial}|F_0|^{2\lambda} v^\lambda \wedge \frac{\eta}{F_0^\ell}$$

vanishes at $\lambda = 0$ if v is a test form that is $\mathcal{O}(|F_0|^\ell)$. However, we may as well assume that F_0 is a monomial in the local coordinates, and therefore this statement follows from the corresponding one-variable statement; that

$$\int_\tau \bar{\partial}|\tau|^{2\lambda} v^\lambda \wedge \frac{\eta(\tau)d\tau}{\tau^\ell}$$

vanishes at $\lambda = 0$ if $\eta = \mathcal{O}(|\tau|^\ell)$. If we write

$$\bar{\partial}|\tau|^{2\lambda} = \lambda|\tau|^{2\lambda} \frac{d\bar{\tau}}{\bar{\tau}}$$

this follows by dominated convergence. $\square$

Proofs of Theorems 1.5 and 1.7. These theorems are proved in much the same way as the corresponding results for holomorphic functions. In fact, if ϕ is a $\bar{\partial}$ -closed smooth form with values in Q , then $(\delta - \bar{\partial})U\phi = \phi - R^f\phi$ so $(\delta - \bar{\partial})U\phi = \phi$ if $R^f\phi = 0$. Following an apparent modification of the procedure in (2.5) we get a $\bar{\partial}$ -closed current ψ with values in E such that $f\psi = \phi$. However, if (1.2) holds, then it is not hard to verify that $U\phi$ is locally integrable; in fact in a resolution as above $U\phi$ is then bounded, in particular it is locally integrable, and therefore $U\phi$ is locally integrable in X . Therefore, we can get a locally integrable solution ψ . $\square$

For further reference we state the following proposition that shows that the principal term R_p^f of the current R^f (where $p = \text{codim } Z$) is robust in certain sense. It is proved precisely as Theorem 2.1 in [5].

Proposition 5.2. *Let f be a generically surjective morphism, let R^f be the associated residue current, and let $p = \text{codim } Z$. Assume that h is a holomorphic section of some line bundle such that $\{h = 0\} \cap Z$ has codimension $p+1$. If ϕ is a holomorphic section such that $R_p^f\phi = 0$ in $X \setminus \{h = 0\}$, then $R_p^f\phi = 0$ in X .*

6. POLYNOMIAL MAPPINGS

Let $L^s = \mathcal{O}(s)$ be the line bundle over $\mathbb{P}^n$ whose sections are represented by s -homogeneous functions in $z = (z_0, \dots, z_n)$. If ϕ is a section of L^s its natural norm is

$$\|\phi(z)\| = \frac{|\phi(z)|}{|z|^s}.$$

Let $E_j, j = 1, \dots, m$, be trivial line bundles over $\mathbb{P}^n$ with basis elements e_j . If f^j are the d_j -homogenizations of the columns of polynomials P^j as in Theorem 1.8, then $f_k = \sum_1^m f_k^j e_j^*$ are sections of

$$E^* = E_1^* \otimes L^{d_1} \oplus \dots \oplus E_m^* \otimes L^{d_m}.$$

Observe that the section $F = f_1 \dots \wedge f_r$ of $\Lambda^r E^*$ can be written

$$F = \sum_{|I|=r}^I F_I e_{I_1}^* \wedge \dots \wedge e_{I_r}^*,$$

where $F_I = \det(f_k^{I_j})$. Thus

$$\|F(z)\|^2 = \frac{\sum_{|I|=r}' |F_I(z)|^2}{|z|^{2 \sum_1^r d_{I_j}}}.$$

If we write this expression in the affine coordinates z' we get precisely the left hand side of (1.4). Let Q be the trivial bundle $\mathbb{C}^r \rightarrow \mathbb{P}^n$. Then f defines a morphism $f: E \rightarrow Q$ such that

$$\psi = (\psi_1, \dots, \psi_m) \mapsto \sum_1^m f^j \psi_j$$

where f^j are the homogenizations of the given (columns) of polynomials P_j of degrees d_j . We can now prove Theorem 1.8.

Proof of Theorem 1.8. Let

$$\dots \rightarrow E_2 \rightarrow E \rightarrow \mathbb{C}^r \rightarrow 0.$$

be the induced complex defined in Section 2. We can take tensor products with L^ρ and get the complex

$$\dots \rightarrow E_2 \otimes L^\rho \rightarrow E \otimes L^\rho \rightarrow \mathbb{C}^r \otimes L^\rho \rightarrow 0.$$

Let U and R^f be the corresponding currents from Section 5 with respect to the natural Hermitian metrics of E and Q . If ϕ is a section of $Q \otimes L^\rho$, and $R^f \phi = 0$, then $v = U\phi$ solves the equations

$$(\delta - \bar{\partial})(v_1 + v_2 + \dots + v_{\min(n+1, m-r+1)}) = \phi.$$

In order to get a holomorphic solution ψ to $f\psi = \phi$ we have to solve all the equations $\bar{\partial}w_k = v_k - \delta v_{k+1}$. Notice that $v_k - \delta v_{k+1}$ is a $(0, k-1)$ -current with values in $E_k \otimes L^\rho$. It is well-known that $H^{0,k}(\mathbb{P}^n, L^\nu) = 0$ for all ν if $1 \leq k \leq n-1$, whereas $H^{0,n}(\mathbb{P}^n, L^\nu) = 0$ (if and only) if $\nu \geq -n$. If $m-r+1 \leq n$ there is therefore no problems at all, and

the only possible obstruction may appear when $k = n + 1$. Notice that v_{n+1} takes values in

$$E_{n+1} \otimes L^\rho = \Lambda^{r+n} E \otimes S^{n-1} Q^* \otimes Q^* \otimes L^\rho.$$

Since Q is trivial, E_{n+1} is a direct sum of line bundles

$$L^{\rho - (d_{I_1} + \dots + d_{I_{r+n}})},$$

where I is an increasing multiindex. Therefore the crucial $\bar{\partial}$ -equation is solvable if $\rho - (d_1 + \dots + d_{n+r}) \geq -n$. Finally we express the relation $\sum f^j \psi_j = f\psi = \phi$ in affine coordinates and get the desired polynomials Q_j as $Q_j(z') = \psi_j(1, z')$. $\square$

Proof of Proposition 1.12. If we have m' polynomials P'_ν of degrees at most d' with no common zeros in $\mathbb{C}^n$, then it is proved in [16] that

$$\frac{|P_\nu(z')|^2}{(1 + |z'|^2)^{d'}} \geq C \frac{1}{(1 + |z'|^2)^M},$$

where $M = (d')^{\min(n, m')}$, provided that $d' \geq 3$. We have $m!/(m-r)!r!$ polynomials $P_I = \det(P_{I_j}^k)$ of degrees $d' = rd$ and hence the proposition follows. $\square$

Proof of Theorem 1.13. Since $\phi = f\psi$ is solvable in $\mathbb{C}^n$ and $p = m - r + 1$, $R^f \phi = 0$ in $\mathbb{C}^n$ by Theorem 1.2. If we take the section $h = z_0$ of $\mathcal{O}(1)$, then since assumption Z has no irreducible component in the plane at infinity, $\text{codim } Z \cap \{z_0 = 0\} = m - r + 2$. By Proposition 5.2 we therefore have that $R^f \phi = 0$ in $\mathbb{P}^n$. Since (1.3) is fulfilled, the desired solution is given by Theorem 1.8. $\square$

7. ESTIMATES FOR A POINTWISE SURJECTIVE MORPHISM

In this section we indicate that our method can be used to get new quite sharp estimates even when f is pointwise surjective. Let us assume that $E \simeq \mathbb{C}^m$ and $Q \simeq \mathbb{C}^r$ are trivial bundles over a smoothly bounded domain $D = \{\rho < 0\}$ in $\mathbb{C}^n$, and equipped with the trivial metrics. Then a morphism $f: E \rightarrow Q$ is just a matrix of holomorphic functions in D . We assume that $f \in H^\infty(D, \text{Hom}(E, Q))$, and that moreover

$$|f_1 \wedge \dots \wedge f_r| \geq \delta > 0;$$

this means that f is uniformly surjective. Notice that since Q is trivial, $\det Q^*$ is just the trivial line bundle, so $F = f_1 \wedge \dots \wedge f_r$. Let

$$\|\phi\|_{H^p}^p = \sup_{\epsilon > 0} \int_{\rho = -\epsilon} |\phi(z)|^p dS, \quad 0 < p < \infty.$$

It was proved in [2] and [3] that if D admits a plurisubharmonic defining function ρ , and $p \leq 2$, then for any $\phi \in H^p(D, Q)$ there is a $\psi \in H^p(E)$ such that $f\psi = \phi$. Moreover, the norm of ϕ is bounded by a constant times $\log(1/\delta)/\delta^{1+\mu}$, where $\mu = \min(n, m - r)$. To be precise, this sharp estimate is only explicitly given in the case $r = 1$, but it follows

(hopefully) in the general case as well with a similar argument. This result is proved by a combination of the L^2 -methods in [22], the refined L^2 estimate for $\bar{\partial}_b$ introduced in [8], and Wolff type estimates.

The case $p > 2$ and $r = 1$ has been studied by several authors in strictly pseudoconvex domains, e.g., [1] and [6], and a generalization to $r > 1$ is made in [14]. These works are based on integral representation and harmonic analysis. There are also similar results for other spaces of functions, see, e.g., the references in [6]. To show how the ideas in this paper can be applied in a situation like this we present the new result Theorem 7.1 below. It is clear that other known results that are proved by means of the Koszul complex in the case $r = 1$ can be generalized to the case $r > 1$ in an analogous way.

Theorem 7.1. *Let D be a strictly pseudoconvex domain with reasonably smooth (C^3 is enough) boundary and $p < \infty$. For any $\phi \in H^p(D, Q)$ there is a $\psi \in H^p(D, E)$ such that $f\psi = \phi$ and*

$$(7.1) \quad \|\psi\|_{H^p} \leq C_\delta \|\phi\|_{H^p},$$

where

$$C_\delta \leq C(\log(1/\delta))^{\mu/2}/\delta^{1+\mu}, \quad \text{if } \mu = \min(n, m-r) > 1$$

and

$$C_\delta \leq C \log(1/\delta)/\delta^2 \quad \text{if } \min(n, m-r) = 1.$$

and $C_\delta = 1/\delta$ if $\min(n, m-r) = 0$, i.e., $m = r$, and C is a constant that is independent of m .

In the case $r = 1$ this coincides with the result in [6]. Since the proof of this generalization follows the proof in [6] quite closely we just give a sketch and indicate the necessary modifications.

Sketch of proof. With the notation as before we have that $v = u\phi$, where $u = u_1 + \cdots u_{1+\mu}$ and

$$u_{k+1} = (\bar{\partial}\sigma)^{\otimes(k-1)} \otimes \sigma \otimes \bar{\partial}\sigma.$$

Since Q is trivial and ϵ_j is a ON-frame we have, cf., Remark 1, that $S = \sum' \bar{F}_I e_I \otimes \epsilon^*$ of $F = \sum' F_I e_I^* \otimes \epsilon$, so the coefficients in S are anti-holomorphic. Moreover, $\sigma = S/|F|^2$ and

$$\sigma = \left(\sum \delta_{f_j} \otimes \delta_{\epsilon_j} \right) \sigma / k!$$

so that

$$\bar{\partial}\sigma = \pm \left(\sum \delta_{f_j} \otimes \delta_{\epsilon_j} \right) \bar{\partial}\sigma / k!.$$

The norm of forms will be taken with respect to the non-isotropic metric $\Omega = -\rho i \partial \bar{\partial} \log(1/\rho)$; we assume that ρ is a strictly plurisubharmonic defining function. For $k \geq 2$ we can estimate the Carleson norm of $(-\rho)^{\frac{k-2}{2}} |u_{k+1}|$ precisely as in the proof of Proposition 5.2 in [6], with F instead of g and σ instead of γ , and obtain the same estimate $C\delta^{-k-1}(\log(1/\delta))^{k/2}$.

The necessary estimate of u_2 is of Wolff type and involves $\mathbb{L}u_2$ where $\mathbb{L}$ is a smooth $(1,0)$ -vector field. More precisely we need to know that the Carleson norm of $|u_2|^2$ is bounded by C^2 and the Carleson norm of $\sqrt{-\rho}|\mathbb{L}u_2|$ is bounded by C , where $C = \delta^{-2} \log(1/\delta)$. It is easily seen that

$$|u_2| \lesssim \frac{|\partial F|}{|F|^3} \leq \frac{1}{\delta} \frac{|\partial F|}{|F|^2},$$

and the desired Carleson estimate of $|u_2|^2$ now follows from Proposition 5.2 in [6]. For any holomorphic function ψ we have that $\sqrt{-\rho}|\mathbb{L}\psi| \lesssim ||\text{partial}\psi|$. When we compute $\mathbb{L}u_2$ we get either derivatives on the factor $1/|F|^2$ or on the functions f_j in $(\sum \delta_{f_j} \otimes \delta_{e_j})^{r-1}$. Therefore we have that

$$\sqrt{-\rho}|\mathbb{L}u_2| \lesssim \frac{\partial h ||\partial F|}{|F|^3} + \frac{|\partial F|^2}{|F|^4} \lesssim \frac{1}{\delta^2} + \frac{|\partial F|^2}{|F|^4},$$

where h is a holomorphic and bounded. It now follows from Proposition 5.2 in [6] that the Carleson norm is $\lesssim \delta^{-2} \log(1/\delta)$ as wanted. $\square$

Remark 2. If we just assume that ϕ is in H^p with respect to the stronger pointwise norm $\|\phi\|$ instead of $|\phi|$, the same proof would give a solution in H^p provided that we could say that the maximal function of $\|\phi\|$ is in $L^p(\partial D)$. However, we do not know if this is true. $\square$

As mentioned above, a similar result has previously been obtained by Hergoualch, [14], using an idea of Fuhrmann, [13], to reduce to the case $r = 1$. However, by this method one has some loss of precision in the dependence of δ . To see this let us describe this method. Assume that $\phi = \sum \phi_j e_j$ is in $H^p(D, Q)$. Now $F = f_1 \wedge \dots \wedge f_r \in H^\infty(D, \Lambda^r E^*)$ and $|F| \geq \delta$, so by the corresponding result for $r = 1$, for each ϕ_j we can find a section H^j of $H^p(D, \Lambda^r E)$ such that $F \cdot H^j = \phi_j$, $j = 1, \dots, r$, with $\|H^j\|_{H^p} \leq C'_\delta \|\phi\|_{H^p}$. Since the rank of $\Lambda^r E^*$ is $m!/(m-r)!r!$, we get

$$C'_\delta \leq C \frac{(\log(1/\delta))^{\mu'/2}}{\delta^{1+\mu'}}, \quad \mu' = \min(n, m!/(m-r)!r!),$$

if $\mu' > 1$. Since $f_1 \wedge \dots \wedge f_r \cdot H^j = \phi_j$, we also have that $f_j \cdot \psi_j = \phi_j$ if

$$\psi_j = (-1)^{j+1} \delta_{f_r} \dots \delta_{f_{j+1}} \delta_{f_{j-1}} \dots \delta_{f_1} H^j.$$

Now $\psi = \psi_1 + \dots + \psi_r$ solves $f_j \psi = \phi_j$ for each j , i.e., $f \psi = \phi$ as wanted, and since f_j are bounded, we get the same estimate

$$\|\psi\|_{H^p} \leq C'_\delta \|\phi\|_{H^p}.$$

However, $\mu' > \mu$ as soon as $m - r < n$ as in this case it is strictly weaker than (7.1).

Remark 3. It is actually possible to solve the equation $F\Psi = \phi$ above with a sharper estimate, by means of the complex

$$\begin{aligned} \rightarrow \Lambda^{r+2}E \otimes S^2Q^* \otimes \det Q^* \rightarrow \\ \Lambda^{r+1}E \otimes Q^* \otimes \det Q^* \rightarrow \Lambda^rE \otimes \det Q^* \rightarrow \mathbb{C} \rightarrow 0 \end{aligned}$$

with the same mappings as before. Combined with the Fuhrmann trick one can then obtain Theorem 7.1. This complex, and the corresponding residues that appear when f is only generically surjective, will be studied in a forthcoming paper. $\square$

REFERENCES

- [1] E. AMAR: *On the corona problem*, J. of Geom. Anal. **1** (1991).
- [2] M. ANDERSSON: *The H^p corona problem in weakly pseudoconvex domains*, Trans. AMS **342** (1994), 241–155, *The H^2 -corona theorem* Preprint Göteborg (1990).
- [3] M. ANDERSSON: *On the H^p -corona problem*, Bull. Sci. Math., **118** (1994), 287–306.
- [4] M. ANDERSSON: *Residue currents and ideals of holomorphic functions*, Bull. Sci. Math., **128**, (2004), 481–512.
- [5] M. ANDERSSON: *The membership problem for polynomial ideals in terms of residue currents*, Ann. Inst. Fourier (to appear).
- [6] M. ANDERSSON & H. CARLSSON: *Estimates of solutions of the H^p and BMOA corona problem*, Math. Ann. **316**, (2000), 83–102.
- [7] C. BERENSTEIN & A. YGER: *Green currents and analytic continuation*, J. Anal. Math. **75** (1998), 1–50.
- [8] B. BERNDTSSON: *$\bar{\partial}_b$ and Carleson type inequalities*, Complex analysis, II (College Park, Md., 1985–86), 42–54, Lecture Notes in Math., 1276, Springer, Berlin, 1987.
- [9] J. BRIANÇON & H. SKODA: *Sur la clôture intégrale d'un idéal de germes de fonctions holomorphes en un point de $\mathbb{C}^n$* , C. R. Acad. Sci. Paris Sér. A **278** (1974), 949–951.
- [10] J-P DEMAILLY: *Estimations L^2 pour l'opérateur $\bar{\partial}$ d'un fibré vectoriel holomorphe semi-positif au-dessus d'une variété kählérienne complète*, Ann. Sci. École Norm. Sup. (4) **15** (1982), 457–511.
- [11] J-P DEMAILLY: *Complex Analytic and Differential Geometry*, Monograph Grenoble (1997).
- [12] A. DICKENSTEIN & C. SESSA: *Canonical representatives in moderate cohomology*, Invent. Math. **80** (1985), 417–434.
- [13] P. FUHRMANN: *On the corona theorem and its application to spectral problems in Hilbert spaces*, Trans. AMS **132** (1968), 55–66.
- [14] J. HERGOUALCH: *Le probleme de la couronne*, Memoire Bordeaux (2001) *Autour du problème de la couronne* Thesis Bordeaux (2004).
- [15] M. HICKEL: *Solution d'une conjecture de C. Berenstein-A. Yger et invariants de contact à l'infini*, Ann. Inst. Fourier **51** (2001), 707–744.
- [16] J. KOLLÁR: *Sharp effective Nullstellensatz*, J. American Math. Soc. **1** (1988), 963–975.
- [17] F.S. MACAULAY: *The algebraic theory of modular systems*, Cambridge Univ. Press, Cambridge 1916.
- [18] M. NÖTHER: *Über einen Satz aus der Theorie der algebraischen Functionen*, Math. Ann. (1873), 351–359.

- [19] M. PASSARE: *Residues, currents, and their relation to ideals of holomorphic functions*, Math. Scand. **62** (1988), 75–152.
- [20] M. PASSARE & A. TSIKH & A. YGER: *Residue currents of the Bochner-Martinelli type*, Publ. Mat. **44** (2000), 85–117.
- [21] H. SKODA: *Application des techniques L^2 à la théorie des idéaux d'une algèbre de fonctions holomorphes avec poids*, Ann. Sci. École Norm. Sup. **5** (1972), 545–579.
- [22] H. SKODA: *Morphismes surjectifs de fibrés vectoriels semi-positifs*, Ann. Sci. École Norm. Sup. **11** (1978), 577–611.

DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY
AND THE UNIVERSITY OF GÖTEBORG, S-412 96 GÖTEBORG, SWEDEN
E-mail address: `matsa@math.chalmers.se`